This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

IN THE CLAIMS:

Please amend claims 1, 5-8 and 10-12; and cancel claims 13-20; add new claims 21 and 22, as follows:

1. (Currently Amended) A method of making an a tissue adhesive, comprising the steps of:

derivatizing collagen with a functional group;

heating a solution including collagen derivatized with a COO functional group to thereby gelatinize said derivatized collagen;

adding derivatized collagen to said solution, said derivatized collagen being derivatized with a COO functional group; and

heating a composition including said derivatized collagen to thereby increase a concentration of said derivatized collagen in said composition repeating said heating and adding steps until the derivatized collagen concentration in said solution is from 300 mg/ml (30%) up to 800 mg/ml (80%), said added derivatized collagen in said solution being gelatinized after said repeated heating steps.

- 2. (Original) A method in accordance with claim 1, further comprising the step of extracting said collagen from a tissue source prior to said derivatizing step.
- 3. (Original) A method in accordance with claim 2, wherein said tissue source includes an animal tissue.
- 4. (Original) A method in accordance with claim 1, wherein said derivatizing step includes a step of reacting said collagen with 4-mercapto-1,8-naphthalic anhydride.

- 5. (Currently Amended) A method in accordance with claim 14

 1, wherein said derivatizing step further includes a step of reaction with glutaric anhydride.
- 6. (Currently Amended) A method in accordance with claim 1, further comprising additional heating steps to adjust said concentration of said derivatized collagen in said composition further comprising the step of derivatizing said collagen with an SH functional group.
- 7. (Currently Amended) A method in accordance with claim 1, further comprising a step of adding a pH altering material to said derivatized collagen to thereby adjust a pH of said composition solution to be within a desired range of 6.8-7.8.
- 8. (Currently Amended) A method in accordance with claim 7

 1, wherein said desired range is 6.8 7.8 further comprising the step of deaerating said solution.
- 9. (Original) A method in accordance with claim 7, wherein said pH altering material includes NaOH.
- 10. (Currently Amended) A method in accordance with claim ± 8, further comprising the step of:

adding a material to said solution, said material being selected from the group of collagen fibrils, collagen fibers and collagen fiber bundles.

11. (Currently Amended) A method of making an adhesive, comprising the steps of:

derivatizing collagen with a functional group; and increasing a concentration of said derivatized collagen in a

composition in accordance with claim 1, further comprising the step of solidifying said solution.

- 12. (Currently Amended) A method in accordance with claim

 11, further comprising the step of extracting said collagen from a tissue source prior to said derivatizing step sectioning said solidified solution into strips.
 - 13. (Cancelled)
 - 14. (Cancelled)
 - 15. (Cancelled)
 - 16. (Cancelled)
 - 17. (Cancelled)
 - 18. (Cancelled)
 - 19. (Cancelled)
 - 20. (Cancelled)
- 21. (New) A method in accordance with claim 1, wherein said added collagen is dried collagen.
- 22. (New) A method in accordance with claim 1, wherein said added collagen is lyophilized.